

DISTORTIONS OF THE HELICOID

JACOB BERNSTEIN AND CHRISTINE BREINER

ABSTRACT. Colding and Minicozzi have shown that an embedded minimal disk $0 \in \Sigma \subset B_R$ in $\mathbb{R}^3$ with large curvature at 0 looks like a helicoid on the scale of R . Near 0, this can be sharpened: on the scale of $|A|^{-1}(0)$, Σ is close, in a Lipschitz sense, to a piece of a helicoid. We use surfaces constructed by Colding and Minicozzi to see this description cannot hold on the scale R .

In [3, 4, 5, 6], Colding and Minicozzi give a complete description of the structure of embedded minimal disks in a ball in $\mathbb{R}^3$. Roughly speaking, they show that any such surface is either modeled on a plane (i.e. is nearly graphical) or is modeled on a helicoid (i.e. is two multi-valued graphs glued together along an axis). In the latter case, the distortion may be quite large. For instance, in [8], Meeks and Weber “bend” the helicoid; that is, they construct minimal surfaces where the axis is an arbitrary $C^{1,1}$ curve (see Figure 2). A more serious example of distortion is given by Colding and Minicozzi in [2]. There they construct a sequence of minimal disks modeled on the helicoid, but where the ratio between the scales (a measure of the tightness of the spiraling of the multi-graphs) at different points of the axis becomes arbitrarily large (see Figure 1). Note, locally, near points of large curvature, the surface is close to a helicoid, and so the distortions are necessarily global in nature.




FIGURE 1. A cross section of one of Colding and Minicozzi’s examples. Here $R = 1$ and $(0, s)$ is a blow-up pair.




FIGURE 2. A cross section of one of Meeks and Weber's examples, with axis the circle. Here R is the outer scale of a disk and (y, s) is a blow-up pair.

Following [4] we make the meaning of large curvature precise by saying a pair $(y, s) \in \Sigma \times \mathbb{R}^+$ is a (C) *blow-up pair* if $\sup_{B_s \cap \Sigma} |A|^2 \leq 4C^2 s^{-2} = 4|A|^2(y)$ (here C is large and fixed and $\Sigma \subset \mathbb{R}^3$ minimal). For Σ minimal with $\partial\Sigma \subset \partial B_R$ where $(0, s)$ is a blow-up pair, there are two important scales; R the outer scale and s the blow-up scale. The work of Colding and Minicozzi gives a value $0 < \Omega < 1$ so that the component of $\Sigma \cap B_{\Omega R}$ containing 0 consists of two multi-valued graphs glued together (see for instance Lemma 2.5 of [7] for a self-contained explanation). On the other hand, Theorem 1.5 of [1] shows that on the scale of s (provided R/s is large), Σ is bi-Lipschitz to a piece of a helicoid with Lipschitz constant near 1. Using the surfaces constructed in [2] we show that such a result cannot hold on the outer scale and indeed fails to hold on certain smaller scales:

Theorem 0.1. *Given $1 > \Omega, \epsilon > 0$ and $1/2 > \gamma \geq 0$ there exists an embedded minimal disk $0 \in \Sigma$ with $\partial\Sigma \subset \partial B_R$ and $(0, s)$ a blow-up pair so: the component of $B_{\Omega R^{1-\gamma} s^\gamma} \cap \Sigma$ containing 0 is not bi-Lipschitz to a piece of a helicoid with Lipschitz constant in $((1 + \epsilon)^{-1}, 1 + \epsilon)$.*

First, we recall the surfaces constructed in [2]:

Theorem 0.2. *(Theorem 1 of [2]) There is a sequence of compact embedded minimal disks $0 \in \Sigma_i \subset B_1 \subset \mathbb{R}^3$ with $\partial\Sigma_i \subset \partial B_1$ containing the vertical segment $\{(0, 0, t) : |t| \leq 1\} \subset \Sigma_i$ such that the following conditions are satisfied:*

- (1) $\lim_{i \rightarrow \infty} |A_{\Sigma_i}|^2(0) \rightarrow \infty$
- (2) $\sup_{\Sigma_i} |A_{\Sigma_i}|^2 \leq 4|A_{\Sigma_i}|^2(0) = 8a_i^{-4}$ for a sequence $a_i \rightarrow 0$
- (3) $\sup_i \sup_{\Sigma_i \setminus B_\delta} |A_{\Sigma_i}|^2 < K\delta^{-4}$ for all $1 > \delta > 0$ and K a universal constant.
- (4) $\Sigma_i \setminus \{x_3 - \text{axis}\} = \Sigma_{1,i} \cup \Sigma_{2,i}$ for multi-valued graphs $\Sigma_{1,i}$ and $\Sigma_{2,i}$.

Remark 0.3. (2) and (3) are slightly sharper than what is stated in Theorem 1 of [2], but follow easily. (2) follows from the Weierstrass data (see Equation (2.3) of [2]). This also gives (3) near the axis, whereas away from the axis use (4) and Heinz's curvature estimates.

Next introduce some notation. For a surface Σ (with a smooth metric) we denote intrinsic balls by $\mathcal{B}_s^\Sigma$ and define the (*intrinsic*) *density ratio* at a point p as: $\theta_s(p, \Sigma) = (\pi s^2)^{-1} \text{Area}(\mathcal{B}_s^\Sigma(p))$. When Σ is immersed in $\mathbb{R}^3$ and has the induced metric, $\theta_s(p, \Sigma) \leq \Theta_s(p, \Sigma) = (\pi s^2)^{-1} \text{Area}(B_s(p) \cap \Sigma)$, the usual (extrinsic) density ratio. Importantly, the intrinsic density ratio is well-behaved under bi-Lipschitz maps. Indeed, if $f : \Sigma \rightarrow \Sigma'$ is injective and with $\alpha^{-1} < \text{Lip } f < \alpha$, then:

$$(0.1) \quad \alpha^{-4} \theta_{\alpha^{-1}s}(p, \Sigma) \leq \theta_s(f(p), \Sigma') \leq \alpha^4 \theta_{\alpha s}(p, \Sigma).$$

This follows from the inclusion, $\mathcal{B}_{\alpha^{-1}s}^\Sigma(f^{-1}(p)) \subset f^{-1}(\mathcal{B}_s^{\Sigma'}(p))$ and the behavior of area under Lipschitz maps, $\text{Area}(f^{-1}(\mathcal{B}_s^{\Sigma'}(p))) \leq (\text{Lip } f^{-1})^2 \text{Area}(\mathcal{B}_s^{\Sigma'}(p))$.

Note that by standard area estimates for minimal graphs, if $\Sigma \cap B_s(p)$ is a minimal graph then $\theta_s(p, \Sigma) \leq 2$. In contrast, for a point near the axis of a helicoid, for large s the density ratio is large. Thus, in a helicoid the density ratio for a fixed, large s measures, in a rough sense, the distance to the axis. More generally, this holds near blow-up pairs of embedded minimal disks:


Lemma 0.4. *Given $D > 0$ there exists $R > 1$ so: If $0 \in \Sigma \subset B_{2Rs}$ is an embedded minimal disk with $\partial\Sigma \subset \partial B_{2Rs}$ and $(0, s)$ a blow-up pair then $\theta_{Rs}(0, \Sigma) \geq D$.*

Proof. We proceed by contradiction, that is suppose there were a $D > 0$ and embedded minimal disks $0 \in \Sigma_i$ with $\partial\Sigma_i \subset \partial B_{2R_i s}$ with $R_i \rightarrow \infty$ and $(0, s)$ a blow-up pair so that $\theta_{R_i s}(0, \Sigma_i) \leq D$. The chord-arc bounds of [7] imply there is a $1 > \gamma > 0$ so $\mathcal{B}_{R_i s}^{\Sigma_i}(0) \supset \Sigma_i \cap B_{\gamma R_i s}$. Hence, the intrinsic density ratio bounds the extrinsic density ratio, i.e. $D \geq \theta_{R_i s}(p, \Sigma_i) \geq \gamma^2 \Theta_{\gamma R_i s}(p, \Sigma_i)$. Then, by a result of Schoen and Simon [9] there is a constant $K = K(D\gamma^{-2})$, so $|A_{\Sigma_i}|^2(0) \leq K(\gamma R_i s)^{-2}$. But for R_i very large this contradicts that $(0, s)$ is a blow-up pair for all Σ_i . $\square$

Remark 0.5. Note that the above does not depend on the strength of chord-arc bounds. In fact, it is also an immediate consequence of the fact that intrinsic area bounds on a disk give total curvature bounds. In turn, the total curvature bounds again yield uniform curvature bounds. See Section 1 of [4] for more detail.

To produce our counterexample, we exploit the fact that two points on a helicoid that are equally far from the axis must have the same density ratio. Assuming the existence of a Lipschitz map between our surface Σ and a helicoid, we get a contradiction by comparing the densities for two appropriately chosen points that map to points equally far from the axis of the helicoid.

Proof. (of Theorem 0.1) Fix $1 > \Omega, \epsilon > 0$ and $1/2 > \gamma \geq 0$ and set $\alpha = 1 + \epsilon$. Let Σ_i be the surfaces of Theorem 0.2; we claim for i large, Σ_i will be the desired example. Suppose this was not the case. Setting $s_i = C a_i^2 / \sqrt{2}$, where a_i is as in (2) and C is the blow-up constant, one has $(0, s_i)$ is a blow-up pair in Σ_i , since $\sup_{\Sigma_i \cap B_{s_i}} |A_{\Sigma_i}|^2 \leq 8 a_i^{-4} = 4 C^2 s_i^{-2} = 4 |A_{\Sigma_i}|^2(0)$, moreover, $s_i \rightarrow 0$. Hence, with $R_i = \Omega s_i^\gamma < 1$, the component of $B_{R_i} \cap \Sigma_i$ containing 0, Σ'_i , is bi-Lipschitz to a piece of a helicoid with Lipschitz constant in (α^{-1}, α) . That is, there are subsets Γ_i of helicoids and diffeomorphisms $f_i : \Sigma'_i \rightarrow \Gamma_i$ with $\text{Lip } f_i \in (\alpha^{-1}, \alpha)$.

FIGURE 3. Finding u_i

We now begin the density comparison. First, Lemma 0.4 implies there is a constant $r > 0$ so for i large $\theta_{rs_i}(0, \Sigma'_i) \geq 4\alpha^8$ and thus by (0.1) $\theta_{\alpha rs_i}(f_i(0), \Gamma_i) \geq 4\alpha^4$. We proceed to find a point with small density on Σ_i that maps to a point on Γ_i equally far from the axis as $f_i(0)$ (which has large density).

Let U_i be the (interior) of the component of $B_{1/2R_i} \cap \Sigma_i$ containing 0. Note for i large enough, as $s_i/R_i \rightarrow 0$, the distance between ∂U_i and $\partial \Sigma'_i$ is greater than $4\alpha^2 rs_i$. Similarly, for $p \in \partial U_i$ for i large, $p' \in \mathcal{B}_{4\alpha^2 rs_i}^{\Sigma'_i}(p)$ implies $|p'| \geq \frac{1}{4}R_i$. Hence, property (3) gives that $|A_{\Sigma'_i}|^2(p') \leq K' s_i^{-4\gamma}$. Thus, for i sufficiently large $\mathcal{B}_{\alpha^2 rs_i}(p)$ is a graph and so $\theta_{\alpha^2 rs_i}(p, \Sigma'_i) \leq 2$. Pick $u_i \in \partial f(U_i)$ at the same distance to the axis as $f_i(0)$ and so the density ratio is the same at both points (see Figure 3). As $f_i(U_i)$ is an open subset of Γ_i containing $f_i(0)$, $p_i = f_i^{-1}(u_i) \in \partial U_i$. Notice that $\theta_{\alpha rs_i}(u_i, \Gamma_i) = \theta_{\alpha rs_i}(f_i(0), \Gamma_i) \geq 4\alpha^4$ so $2\alpha^4 \geq \alpha^4 \theta_{\alpha^2 rs_i}(p_i, \Sigma'_i) \geq 4\alpha^4$. $\square$

REFERENCES

1. J. Bernstein and C. Breiner, *Helicoid-like minimal disks and uniqueness*, Preprint.
2. T. H. Colding and W. P. Minicozzi II, *Embedded minimal disks: Proper versus nonproper-Global versus local*, Trans. AMS **356** (2003), no. 1, 283–289.
3. ———, *The space of embedded minimal surfaces of fixed genus in a 3-manifold I; Estimates off the axis for disks*, Ann. of Math. (2) **160** (2004), no. 1, 27–68.
4. ———, *The space of embedded minimal surfaces of fixed genus in a 3-manifold II; Multi-valued graphs in disks*, Ann. of Math. (2) **160** (2004), no. 1, 69–92.
5. ———, *The space of embedded minimal surfaces of fixed genus in a 3-manifold III; Planar domains*, Ann. of Math. (2) **160** (2004), no. 2, 523–572.
6. ———, *The space of embedded minimal surfaces of fixed genus in a 3-manifold IV; Locally simply connected*, Ann. of Math. (2) **160** (2004), no. 2, 573–615.
7. ———, *The Calabi-Yau conjectures for embedded surfaces*, Ann. of Math. **167** (2008), no. 1, 211–243.
8. W.H. Meeks and M. Weber, *Bending the helicoid*, Mathematische Annalen **339** (2007), no. 4, 783–798.
9. R. Schoen and L. Simon, *Regularity of simply connected surfaces with quasiconformal Gauss map*, Seminar on minimal submanifolds, Ed. E. Bombieri, Ann. of Math. Studies **103**, 127–145.